

Allocasuarina verticillata Drooping She-oak

TAXONOMY

Division Angiosperm
Subclass Dicotyledonae
Family Casuarinaceae

Previous Taxonomic Names

Casuarina verticillata Lam. (ANH et al 2005)
Casuarina stricta Aiton (ANH et al 2005)
Casuarina quadrivalvis Labill. (ANH et al 2005)

Taxonomic Identification Number 28220 (ANH et al 2005)

Taxonomic Status

Long-lived perennial

Common Names

Drooping She-oak, Coast She-oak, Hilloak; also various spelling variations such as She Oak, Sheoak, She oak, She-oak (ANBG n.d.).

MORPHOLOGY

Tree 4-10 m tall with separate male and female plants (dioecious).
Bark furrowed, branchlets drooping, grey-green, to 40 cm long. Internodes 10-40 mm long, 0.7-1.5 mm diameter, broadened at end near visible teeth.
Male plants with yellow-brown flower spikes, 3-12 cm long, 2-4 whorls per centimetre.
Female flowers red, globular. Large, cylindrical to barrel-shaped cones, to 20-50 m long, 17-30 mm in diameter, which usually have pointed valves (Costermans 1983; Walsh & Entwistle 1996).

SUBSPECIES

None

HYBRIDS

None known

SIMILAR SPECIES

Allocasuarina luehmannii (Buloke) - readily distinguishable by examining internode shape and length, length of male flower spikes, and the size and shape of cones (Walsh & Entwistle 1996).

GEOGRAPHIC RANGE

Found throughout Victoria, except in the Mallee and Gippsland and Alpine highlands. Prefers well-drained soils such as rocky coastal areas, basalt plains and hills and on dry, rocky highland hills and ridges. Also SA, NSW and TAS (Walsh & Entwistle 1996).

Lyons, Brooks and Craig (1974) found that in the Coolac Serpentinite belt in NSW where *A. verticillata* is one of the typical species, a highly significant plant-soil relationship existed between *A. verticillata* and zinc. They also noted that the distribution of *A. verticillata* in that region appeared to be controlled by higher potassium and nickel values in the soil.

BIOREGIONS

Central Victorian Uplands Otway Plain Otway Ranges
Warrnambool Plain Victorian Volcanic Plain

PLANT COMMUNITIES

In Corangamite, *A. verticillata* is associated with grasslands and grassy woodlands on the volcanic plains, scrubs and woodlands along the coast, wet forests in the Otway ranges and dry forest ecosystems especially around central Victoria. Known to form pure woodland stands.

FRAGMENTATION

Fragmentation is recent, occurring since European settlement.

RELEVANT HISTORY & RESEARCH

Moncur et al (1997) states that there has been little studied regarding the reproductive biology of the Casuarinaceae family. No studies regarding floral biology with regard to management of seed production or seed orchards and controlled crosses were completed prior to the 1997 study by Moncur, Boland and Harbard.

Several studies on cockatoos highlight the importance of *Allocasuarina verticillata* as a food source. Schodde *et al* (1993) noted that Drooping Sheoke was a significant species for a variety of parrots that feed on the fruit through out the year. Cleland and Sims (1968) and Cameron (2005) reported that the Glossy Black-cockatoo which has a very specialised diet, feeds almost entirely on *Allocasuarina* and *Casuarina* seeds.

Current research through the Corangamite Seed Supply Framework (DPI and GAV) and CSIRO is focused on population genetics of Drooping Sheoke in linear roadside remnants, patches, isolated plants in paddocks and revegetation sites in the Corangamite region. Results from this research will be finalised in December 2006 and will provide clearer guidelines for seed collection and revegetation practises.

POPULATION DENSITY

Small stands are known along roadsides with some large stretches of 100 plants or more. There are also some large stands on private land and public reserves. Most stands across the region are small and isolated.

BREEDING SYSTEMS

FLOWERING

Separate male and female trees. Flowers all year. Male trees have yellow-brown flower-spikes. Female trees have red globular flowers (Walsh & Entwistle 1996).

Bonney (2003) describes a March-December flowering period for the species while Moncur et al (1997) notes flowering during winter months.

A natural stand of 95 trees in Canberra was the subject of a study by Moncur et al (1997). Flowering occurred from May-October (autumn & spring). Most trees in the stand were synchronised from June to mid-October, but once an increase of air temperatures occurred, abundance of flowers was reduced.

Female flowers have red stigmas, which turn brown within two weeks of being pollinated. Stigmas are receptive over their entire length, and are most successfully pollinated from the bottom third of the stigma. Few losses of flowers were observed between fertilisation and maturity (Moncur et al 1997).

POLLEN

Male flowers release pollen over a 3-9 week period depending on air temperature. Pollen germination was highest at 10°C and 15°C and poor at 5°C, 20°C and 25°C. Differences in the levels of pollen viability were recorded, both between individual trees, and for particular trees during different times during the flowering period. Each male flower contains around 650 000 pollen grains. Pollen may have a short life span but is shed over a long time allowing pollination of nearby female trees, depending on wind direction. Male trees are suitably adapted for wind pollination strategies. The total number of viable pollen grains in a given inflorescence was proven to produce 100% pollination, despite the presence of non-viable grains (Moncur et al 1997).

POLLINATION

Obligate out-breeder (Moran 1992; Walsh & Entwistle 1996; Moncur et al 1997).

In the Moncur et al (1997) study, individual female flowers could be receptive for over 12 weeks. The proportion of individual flowers pollinated increased gradually over this period. A number of female trees were receptive 4 weeks prior to the male trees in the same stand shedding their pollen and were being pollinated

from another, more distant stand. Time from pollination to 50% fertilisation, measured by swollen ovaries, took 53-83 days. Pollination is a gradual process which increases over time. These strategies allow a better mix of genes as each flower can receive pollen from a number of males and males can distribute pollen to many females.

POLLINATORS

Wind pollinated (Moran 1992; Moncur et al 1997).

Moncur et al (1997) concluded that any pollination occurring from birds feeding on seeds would be purely accidental, and that the species was pollinated by wind. On warm days pollen clouds were observed, indicating pollen transfer by wind.

SEED

SEED DESCRIPTION

Cylindrical or barrel-shaped woody cones 20-50 mm long and 17-30 mm in diameter. Valves in several rows, often pointed.

Seed 7-12 mm long, very dark brown in colour, with a flat papery wing attached (Walsh & Entwistle 1996). It becomes mucilaginous when wetted (Moncur et al 1997).

Information relating to seed weight and viability varies:

- 130-278 seeds/gram (GAV n.d.) to around 400 seeds/gram (Gowers 1990).
- 127-270 viable seeds/gram with an average seed viability of 58% (Langkamp 1987). Seed from a Canberra population study was found to have an average viability of only 27.4%, with a 25.8% germination rate at 25°C. Cones averaged 93 seeds, with an average seed weight of 0.93g. (Moncur et al 1997).
- 46-70 germinants/gram (GAV n.d.) to 57 germinants/gram at 15°C (Gunn 2001).

SEED CROP

Collect throughout the year, as seeds retained in cones on trees for long periods.

Collect cones closest to the trunk. Mature cones will be grey-brown in colour, but ensure valves remain closed. Immature fruits are tan brown, and immature seed is pale cream to orange in colour, while mature seed is brown (Bonney 2003).

SEED DISPERSAL

Birds (Schodde et al 1993; Moncur et al 1997; Bonney 2003).

Important constant food source for a number of parrots (Moncur et al 1997) including the black cockatoo (Schodde et al 1993).

At no time during Moncur et al's (1997) six month study were any significant quantities of seed dropped by *A. verticillata* trees. This confirms Turnbull and Martensz's (1982) findings, and supports their suggestion that fire may be required for seed to be released (Moncur et al 1997).

EXTRACTION & STORAGE

Place collected cones in a warm, dry area and seed will be released in a few days (Bonney 2003).

Seed with a pre-storage viability of 58%, was found to have reduced in viability to 50% after 16 years in storage at 20-25°C (Turnbull & Martensz 1983).

Seed stored at 3-5°C had a 10% germination rate after 5 years, and 83% germination rate after 10 years (Gunn 2001).

PROPAGATION

Can be propagated from fresh seed (Earl et al 2001), although Bonney (2003) advises that best results will be achieved from seed sown between one and eight weeks after collection.

Bonney (2003) states the ideal temperature for germination is 20-25°C, while Gunn (2001) gives the ideal temperature as 15°C. Moncur et al (1997) found that between 10-20°C Drooping Sheoke germinated well. Studies by Turnbull and Martensz (1982) noted that *A. verticillata* from a coastal site at Bermagui germinated well at 25-30°C. This was a much warmer site than the Canberra study.

Pale foliage and poor growth may result unless seedlings are inoculated with soil bacterium (*Franks* spp.) that can be obtained from the roots of parent trees. The bacterium occurs in fleshy and coral-

like root nodules just beneath the soil surface. Dig up small quantity of nodules and grind using mortar and pestle. Dilute and water into seedlings (Earl et al 2001).

Root suckers can be transplanted during winter when soil is moist (Earl et al 2001).

An Ocean Grove study found that moderate shade treatment of 30% of full daylight for nine months between autumn and spring increased the growth rate of *A. verticillata* (Withers 1979a), but prolonged shading decreased the species drought resistance (Withers 1979b). Deep shade treatment of 8% of full daylight was detrimental to growth rates (Withers 1979a) and decreased drought resistance further (Withers 1979b).

TREATMENT OPTIONS

No treatment is required, although inoculation with appropriate mycorrhiza (soil bacterium) may improve results (GAV n.d.; Earl et al 2001). Moncur et al (1997) suggest that stratification should be a standard practice for this species. They found that stratification at 4°C for 4 weeks increased the speed of germination, although it did not affect the total germination rate.

Moncur et al (1997) found that the application of gibberellins (GA₃) had a slight beneficial effect on total germination rates.

GERMINATION TIME

See notes under 'Propagation' in relation to variation in optimal germination temperatures.

Published germination rates and times include:

- Untreated seed took 5 days to reach 50% germination at 25-30°C (Fox et al 1987).
- Untreated seed from Ocean Grove took 35 days at a constant temperature of 26°C to gain near complete germination (Withers 1978).
- Untreated seed should germinate in 2-5 weeks at hot temperatures of up to 50°C (Earl et al 2001).
- Seed from a Canberra site reached 50% of final germination after 14 days, but continued to germinate for 12 weeks. Commencement of germination was delayed by lower temperatures (Moncur et al 1997).

FIELD ESTABLISHMENT

Well suited to direct seeding (Ralph 1994; Bonney 2003).

Sow into lightly tilled soils from late winter to early spring through a machine boot and press in (Bonney 2003).


Natural regeneration is from root suckers and seed, but it will also coppice to some extent. Highly palatable to livestock and rabbits, so generally requires protection to establish (Earl et al 2001).

SEED COLLECTION RANGE - *Allocasuarina verticillata*

Intermediate - within which, collection can be extended to formally contiguous remnants

Coastal populations of *Allocasuarina verticillata* vary in form to inland populations. These areas differ markedly in rainfall and geology. Separate collections should be maintained for inland and coastal areas.

Collections can extend to formally contiguous populations. Collect from a large number of female plants (30-50 plants) where possible. Collect from populations with good numbers of both female and male plants. Branches should not be lopped as this impacts dramatically on future seed set. The cones can be hand picked. Collect greying cones as these are more mature than brown cones.



MAP: *Allocasuarina verticillata* broad distribution

DATA SOURCE: Flora Information System, Biodiversity and Natural Resources, DSE –May 2005



Broad distribution of remnant patches and scattered trees

REFERENCES

- ANBG n.d., *Australian plant common name database*, Australian National Botanic Gardens, Dept. of Environment & Heritage, retrieved 18 Apr 2006, <<http://www.anbg.gov.au/common.names>>
- ANH, ANGB & ABRS 2005, *Australian plant name index*, Australian National Herbarium, Australian National Botanic Gardens, Australian Biological Research Study, retrieved 18 Apr 2006, <<http://www.anbg.gov.au/cpbr/databases/apni.html>>
- Bonney, N 2003 (2nd edition), *What seed is that?*, Neville Bonney, Tantanoola, SA.
- Cameron, M (2005) Group size and feeding rates of Glossy Black-Cockatoos in central New South Wales. *Emu* 105, 299-304
- Costermans, L 1983 (revised edition), *Native trees and shrubs of south-eastern Australia*, New Holland Publishers, Sydney, Australia.
- Earl, G, Stelling F, Titecumb, M & Berwick, S (eds.), 2001, *Revegetation guide for the Goulburn Broken Catchment*, Dept. of Natural Resources & Environment, Melbourne, VIC.
- Gowers, LJ 1990, *Native trees and shrubs of the Ballarat region*, Dept. of Conservation & Environment, Ballarat, VIC.
- Greening Australia Victoria (comp.), n.d., *Indigenous plants for North Central Victoria: a revegetation guide*, Dept. of Natural Resources & Environment, Melbourne, VIC.
- Gunn, B 2001, *Australian Tree Seed Centre: Operations Manual*, CSIRO Forestry and Forestry Products, Canberra, ACT.
- Fox, J, Dixon, B, Monk, D 1987 'Germination in other plant families' in *Germination of Australian Native Plant Seed*, PJ Langkamp (ed), Inkata Press, Melbourne & Sydney.
- Langkamp PJ (ed) 1987, *Germination of Australian Native Plant Seed*, Inkata Press, Melbourne & Sydney.
- Lyons, MT, Brooks, RR & Craig, DC 1974, 'The influence of soil composition on the vegetation of the Coolac Serpentine Belt in New South Wales' in *Journal and Proceedings, Royal Society of New South Wales*, vol.107, pp.67-75.
- Moncur, MW, Boland, DJ & Harbard, JL 1997, 'Aspects of the floral biology of *Allocasuarina verticillata* (Casuarinaceae)', *Australian Journal of Botany*, vol.45, pp.857-869.
- Moran, GF 1992, 'Patterns of genetic diversity in Australian tree species' in *New Forests*, vol.6, pp.49-66.
- Ralph, M 1994 (2nd edition), *Seed collection of Australian native plants : for revegetation, tree planting and direct seeding*, Bushland Horticulture, Fitzroy, VIC.
- Schodde, R, Mason, IJ & Wood, JT 1993, 'Geographic differentiation in the glossy black-cockatoo *Calyptohynchus lathami* (Temminck) and its history', *Emu*, vol.93, pp.156-166.
- Turnbull, JW & Martensz PN 1983, 'Seed production, collection and germination in Casuarinaceae' in *Casuarina ecology management and utilization : proceedings of an international workshop*, Canberra, Australia 17-21 August 1981 (eds. Midgley, S.J., Turnbull, J.W. & Johnston, R.D.), pp.126-132, CSIRO, Melbourne.
- Walsh, NG & Entwistle, TJ (eds.) 1996, *Flora of Victoria*, vol. 3, Inkata Press, Melbourne, VIC.
- Withers, JR 1978, 'Studies on the status of unburnt *Eucalyptus* woodland at Ocean Grove, Victoria. II The differential seedling establishment of *Eucalyptus ovata* Labill. and *Casuarina littoralis* Salisb.' in *Australian Journal of Botany*, vol.26, pp.465-83.
- Withers, JR 1979a, 'Studies on the status of unburnt *Eucalyptus* woodland at Ocean Grove, Victoria. IV The effect of shading on seedling establishment', *Australian Journal of Botany*, vol.27, pp.47-66.
- Withers, JR 1979b, 'Studies on the status of unburnt *Eucalyptus* woodland at Ocean Grove, Victoria. V The interactive effects of droughting and shading on seedlings under competition' in *Australian Journal of Botany*, vol.27, pp.285-300.

ACKNOWLEDGEMENTS

Information compiled for this note series was a result of extensive literature reviews and plant record searches completed by Lucy Nuttal, with assistance from Michelle Butler, Christine Gartlan and Anne Ovington for the Corangamite Seed Supply Framework project.

To contribute to or provide feedback on this note, please email the Corangamite Seed Supply & Revegetation coordinators:

Michelle Butler, DPI michelle.butler@dpi.vic.gov.au
Chris Gartlan, GAV christine.gartlan@dpi.vic.gov.au

DISCLAIMER

The authors do not guarantee that the information is without flaw of any kind or wholly appropriate for your purposes and therefore disclaim all liability for any error, loss or consequence that may arise as a result of you relying on this information.

2006